

Nautiloid cephalopods – a review of their use and potential in biostratigraphy

D.H. EVANS, A.H. KING, K. HISTON & M. CICHOWOLSKI

Abstract: In terms of their use as biostratigraphical tools, nautiloid cephalopods are the poor relations of ammonoids. Nevertheless, in certain situations, they may provide useful biostratigraphical data, particularly where other biostratigraphically valuable taxa are not present; or in certain situations demonstrate a resolution as great as, or greater than ammonoids, trilobites, graptolites or conodonts. Nautiloid cephalopods are of especial value in palaeobiogeographical studies, but their use for this purpose may be hampered by the poor understanding of the stratigraphical ranges of individual taxa. The biostratigraphical value of nautiloid cephalopods is demonstrated here through a number of case studies of Ordovician taxa, combined with a review of their biostratigraphical use in Palaeozoic and Mesozoic successions. These both demonstrate the potential of this group and indicate great scope for further research.

Keywords: Nautiloids, cephalopods, biostratigraphy, Palaeozoic, Mesozoic

Introduction

The widespread stratigraphical and geographical occurrence of nautiloid cephalopods, particularly, but not exclusively during the Lower Palaeozoic is documented in many substantial monographic works (e.g. HALL 1847 [Ordovician and Silurian]; BARRANDE 1865-1877 [Ordovician to Devonian]; BLAKE 1882 [Ordovician and Silurian]; FOORD 1897-1903 [Carboniferous]; FOERSTE 1932, 1933 [Ordovician]; FLOWER 1946 [Ordovician]; KUMMEL 1953 [Triassic]; MILLER & YOUNGQUIST 1949 [Permian]; STURGEON et al. 1997 [Carboniferous]; ZHURAVLEVA 1974 [Devonian]). Unlike ammonoids, which have been utilised as stratigraphical tools since the early nineteenth century (e.g. SMITH 1816; OPPEL 1865; BUCKMAN 1898; HOUSE 1978; CALLOMAN et al. 1989; KORN 1996; KLUG 2002; BECKER 2000; PAGE 2009), and despite the great variation in conch morphology that ‘nautiloids’ display, examples of their use in the development of biostratigraphical schemes are relatively scarce. There are several possible reasons for this:

1. Biostratigraphical markers: There are a range of other rapidly evolving and widely occurring taxa such as graptolites, ammonoids, conodonts and foraminifera that are of proven value, and may occur in great abundance, facilitating correlations using samples extracted from cores.

2. Similarity of morphology: Many of the groups comprising these cephalopods cannot be determined

taxonomically without investigation of the internal morphology of the phragmocone due to the similarities of the external features, thus entailing the preparation of polished or thin sections for study, requiring more effort, and possibly resulting in more equivocal data than might be gained from the preparation of (for example) a conodont sample.

3. Taphonomy: Amongst the many forms with orthoconic shells, there is great potential for post-mortem – pre-burial breakage and preferential removal of parts of the phragmocone to bring about the selective preservation of parts of the shell on which the diagnosis of a particular taxon may be based. Taphonomic studies of the post-mortem behaviour and deposition of nautiloid cephalopod shells (REYMENT 1958; BOSTON & MAPES 1991; HEWITT & WESTERMANN 1996; HISTON 2012a) deal with many aspects of their preservation. If there is more than one mode of preservation for a particular taxon, and particularly if these occur at different horizons, there is potential for mistakenly splitting that taxon and failing to recognise its full stratigraphical range. For example, *Polymeres demeterum* MURCHISON from the Floian of England and Wales occurs at several different horizons where the preserved remnants of conch represent either adoral, medial or apical portions, whilst their preservation may take the form of internal and external moulds, lacking original shell, or original shell may have been replaced by pyrite followed by further replacement with limonite. Without careful study of this material and an understanding of its taphonomy,

it would be easy to determine these as separate taxa and set up spurious ranges for them.

4. Fidelity of palaeogeographical distribution: An important factor in this discussion is whether the nautiloid shell could float after death of the animal as then the original palaeoenvironmental setting and any hypothesis related to facies and assemblages would be prone to error. This has led to the use of nautiloids as reliable biostratigraphic markers and precise indicators of palaeoenvironment being treated with doubt or completely dismissed in holostratigraphical studies. HEWITT & WESTERMANN (1996) concluded that post-mortem buoyancy of nautiloid shells was limited as most individuals lived on or near the seafloor and would sink soon after death rather than float to the surface as the cameral chambers flooded. Therefore, it is considered that nautiloid shells were deposited on the seafloor shortly after the death of the organism and would have remained buoyant for only a short period, if at all (KRÖGER et al. 2009; KLUG et al. 2010; HISTON 2012b).

As a probable consequence, examples of the direct application of nautiloid cephalopods as biostratigraphical tools are relatively few. However, this is principally due to the fact that nautiloid biostratigraphical zones/schemes have never been attempted, not as a result of failure in their application. More often, biostratigraphical studies of these cephalopods are markedly descriptive, and may largely represent an extension of a monographic study (e.g. FREY 1995; STURGEON et al. 1997; KRÖGER 2008a; EVANS 2005, 2011), or the provision of known ranges for a particular region/locale (e.g. WILSON 1961; CATALANI in SLOANE 1987). Rousseau FLOWER was probably one of the main exponents of the use of these cephalopods as biostratigraphical tools, and used his knowledge of their distribution, particularly in North America, to review the biostratigraphy of various successions of Lower, Middle and Upper Ordovician age and also as a tool to better understand the evolution of this group of cephalopods (see FLOWER 1976; 1985 for summaries). In one instance, FLOWER (1964) used his knowledge of the stratigraphical distribution of Canadian (= Lower Ordovician) cephalopods of North America to date some of the larger clasts present in the Levis Conglomerate of Quebec, demonstrating that the boulders originated from carbonates ranging (in modern terminology) from Skullrockian to Blackhillsian in age, and indicating a source or sources further shoreward on the Laurentian platform. In this case the general lack of other macrofossils made these cephalopods a key to the age determination of this material, and although today, age would probably be determined by conodont sampling, the value of these cephalopods in assisting age determination in situations where other macrofossils are

relatively rare, as in the Durness Group of Northwest Scotland (EVANS 2011) should not be underestimated.

The study of the distribution of nautiloid cephalopods for palaeobiogeography and as tools for palaeogeographical reconstruction was strongly proposed by CRICK in the 1980's and 1990's (CRICK 1980, 1988, 1990, 1993). He put forward valid arguments supported by a sound database of systematic works to support his hypotheses that these faunas are particularly sensitive to distance or water depth separating land-masses and to fluctuations in sea level. As CRICK pointed out faunas should be described using precise systematic criteria within a strictly controlled biostratigraphic framework in order to fully exploit their potential. When CRICK published his major contributions this was not always the case and many systematic studies were lacking a precise stratigraphic context even at series level, many taxa being referred to as "Upper Ordovician", "lower Silurian" etc. These seminal works by CRICK gave impetus to a broad array of studies over the last twenty years by other authors (e.g. CICHOWOLSKI, EVANS, FREY, HERWIG & POSENATO, HISTON, HOLLAND, KING, KLUG, KRÖGER, MAPES, and NIKO among others: see reference list for details) on these faunas from a variety of geographic locations ranging stratigraphically from the Ordovician to the Triassic that have shown without doubt that nautiloid cephalopods are indeed reliable palaeobiogeographical indicators. Nautiloid studies are now much more accurate in this respect when based on newly collected material, however, material being redescribed from historical collections is still a major problem and as CRICK suggested, is often only of use from a taxonomic point of view.

More recent studies of nautiloid cephalopods may recognise the difficulties that can arise when attempting to use these organisms as biostratigraphical tools and make use of new collections sampled within a well-constrained biostratigraphical framework. Studies of the Silurian cephalopods from Europe in particular (e.g. HISTON 2012a; HISTON et al. 2010; MANDA et al. 2009) make use of such a well-constrained biostratigraphical framework to accurately constrain the range of taxa occurring within the succession, and contributes, in combination with previous studies of other European faunas (see sections below for references) to their use in biostratigraphical correlation as well as a palaeobiogeographical tool within the framework of Northern Gondwana. Such an approach, which recognises that the stratigraphical distribution of these cephalopods should be assessed within a background of other biostratigraphical constraints are more likely to yield useful results in terms of developing their biostratigraphical value.

Examples from recent studies

There are a number of situations where the real or potential biostratigraphical value of these cephalopods may be of significance. Where other biostratigraphically critical fossil biota may be relatively rare, as is the situation along parts of the Laurentian margin during the Early Ordovician (above), areas of the North Gondwana margin where Silurian cephalopod limestones dominate, or in parts of the Early Jurassic successions of the North Somerset coast (United Kingdom) described further below. Where these cephalopods are particularly abundant, as in parts of the Ordovician successions of Scandinavia and South China, some groups, such as the lituitids (see below), underwent a rapid evolution, and may be distinguished as a sequence of distinct taxa that in some cases can be recognised on both palaeoplates, and may potentially be used as correlative tools. Studies by MANDA (2009) of the phragmocerids from the classic Silurian deposits of the Prague Basin also demonstrate this aspect. Use of Silurian nautiloid assemblages by GNOLI (1990) and MANDA & KRIZ (2006) highlight the potential for correlation. Significant, and/or relatively rapid evolutionary events during the history of these cephalopods may also have some degree of biostratigraphical value, and examples from the early history of the Orthoceratoidea and the Eothinoceratidae are described here. Finally, where the stratigraphical distribution of cephalopods are well constrained, whilst they may not provide the primary evidence on which to found a biostratigraphical scheme, may provide a significant contribution to more comprehensive, holostratigraphical schemes.

The examples provided below have a strong Ordovician and lower Palaeozoic bias, reflecting the research interests of the authors. As such, they provide an indication of the potential for the use of nautiloid cephalopods as biostratigraphical tools, and indicate, together with the Mesozoic examples that these cephalopods can be used in a biostratigraphical context.

Ordovician

1. Orthoceratoidea. Research carried out during the past decade has pushed back the origins of the Class Orthoceratoidea well into the late Tremadocian (KRÖGER 2008; KRÖGER & EVANS 2011) whilst indicating that diversification of these cephalopods was already taking place in the Tremadocian (KRÖGER & EVANS 2011) and increased in the early Floian (EVANS 2005, 2011) as demonstrated by the extended ranges of several Laurentian taxa (EVANS 2011, and Fig. 1 herein).

The proposed revision of the position of the Stairian-Tulean boundary to approximately coincide with

the Tremadoc-Floian boundary based on the stratigraphical and palaeogeographical distribution of pelagic trilobites (ADRAIN 2011) indicates that the delay between the appearance of the Orthoceratoidea in high latitude sites on the west Gondwana margin and at low latitudes (NW Scotland, Newfoundland, SW United States) is significantly less than previously thought. Such a rapid dispersion (if that is what we are observing) of this group of cephalopods would appear to be in accord with a move to a more pelagic habitat (KRÖGER et al. 2009) and would suggest that the adaptations noted by KRÖGER (2005); particularly the appearance of a small subspherical protoconch that may have facilitated the rapid dispersal of large numbers of offspring, took place very early in the history of the group and is likely to be present in all the taxa referred to in Fig. 1, despite the lack of any evidence of this morphology in most of these taxa as a consequence of the rarity of preservation of the apical portion of the phragmocone.

At present, with the exception of *Slemmestadoceras attavus* (BROEGGER), the earliest known members of the Orthoceratoidea are represented by taxa possessing marginal siphuncles. These may be assigned to *Bactroceras* HOLM, or closely related genera, all of which so far have shown no evidence of the presence of cameral deposits, and only *Thoraloceras bactroceratoides* KRÖGER & EVANS possesses endosiphuncular deposits that may be interpreted as a conical lining that is thicker at the septal necks than along the rest of the siphonal segment. Given that this taxon is known only from very fragmentary material and it is not known whether cameral deposits might have been present, some doubt will remain regarding its assignment to the Troedsonellidae (Dissidocerida).

Uncertainty regarding an appropriate assignment for *Thoraloceras*, together with the presence of *Bactroceras* in the same assemblage, combined with a possible *Cochlioceras* that is marginally younger means that it is not possible to resolve the relationships between these taxa without additional data. A cladistic analysis (KRÖGER 2008b) generated over one hundred similarly parsimonious trees, although it should be noted that characters including the presence/absence and nature of endosiphuncular and cameral deposits were not included in the analysis. Comparison of the two trees figured by KRÖGER (2008b, fig. 3) illustrates the problem. Nevertheless, if this group of taxa achieved their wide distribution through (at least in part) the innovation of the small spherical protoconch, then this could provide the character that unites the Orthoceratoidea. This implies that the late Tremadocian and the Floian was a period of rapid radiation for the Orthoceratoidea, much of the documentation of which remains to be discovered.

2. Eothinoceratidae and Bathmoceratidae. Although the families Eothinoceratidae and Bathmoceratidae achieved a distribution across several palaeocontinents during the Floian and Darriwilian, the oldest member of the Eothinoceratidae known is considered to belong to *Saloceras sericeum* (SALTER), represented by *S. cf. sericeum* from the Floresta Formation of the Cordillera Oriental of northwest Argentina (CICHOWLSKI & VACCARI 2011), and by the lost type material of *Cyrtoceras praecox* SALTER, here considered likely to belong to *S. sericeum* (see discussion in EVANS 2005, p. 67) that originated from the Dol-cyn-afon Formation (*tennelus* graptolite biozone) of North Wales. As the material described by CICHOWLSKI & VACCARI (2011) came from the *deltifer* conodont biozone, these two records are of broadly the same age, and appear to mark the first appearance of cephalopods in high latitudinal Gondwana.

cations around Gondwana (EVANS 2007; CICHOWOLSKI & VACCARI 2011; KRÖGER & EVANS 2011) and certainly underwent diversification as indicated by the assemblages from the Montagne Noire (KRÖGER & EVANS 2011). *Proterocameroceras contrarium* TEICHERT & GLENISTER from the Emanuel Limestone of northwest Australia may belong to *Saloceras* (EVANS 2005, p. 11) or represent a new genus of eoithinoceratid. *Saloceras* ranges into the mid and late Floian in the Welsh Basin (EVANS 2005) and the Central Andean Basin (CICHOWOLSKI unpublished data).

The earliest records of *Eothinoceras* are from the latest Tremadoc and earliest Floian of low latitude Gondwana (Western Australia [TEICHERT & GLENISTER 1954]) and Laurentia (ULRICH et al. 1944; KRÖGER & LANDING 2008). The occurrence of the genus in the Rochdale Formation of New York State suggests that it may be slightly older than *Protothinoceras* CHEN & TEICHERT from the early Floian Liangchiashan Formation of Hebei Province, North China and regarded as the ancestor of *Eothinoceras* by CHEN & TEICHERT (1987, text-

			Avalonia	Montagne Noire	Morocco	Perunica	Andean Basin	Baltica	Southern China	Laurentia	Australia	North China
Lower Ordovician	Middle Ordovician											
	Dapingian	Darriwilian										
	Floian											
Tremadocian	1a-1d											
	2a-2c											

Fig. 2: Worldwide stratigraphical distribution of members of the Eothinoceratidae and Bathmoceratidae during the Early and Middle Ordovician.

fig. 3). Nevertheless, the fauna of the Liangchiashan Formation clearly records what may have been a short-lived proliferation of genera that could have arisen from *Eothinoceras* and probably gave rise to the Cyrtocerinidae through *Tangshanoceras* CHEN (CHEN & TEICHERT 1987). The appearance of *Eothinoceras* at distant locations during the latest Tremadoc and early Floian is difficult to explain unless it is accounted for by a longer history of the genus or a capacity for rapid dispersal. *Eothinoceras renatae* CECIONI & FLOWER from the Floian of Bolivia, and *E. marchaense* BALASHOV from the Floian of Siberia provide further evidence of this wide, if sporadic, distribution of this genus during the Floian.

The occurrence of *Eothinoceras* and *Saloceras? contrarium* (TEICHERT & GLENISTER) in the Canning Basin of Western Australia, combined with the occurrence of *Bathmocerases australe* (TEICHERT) from the late Floian-early Dapinginian Horn Valley Siltstone (COOPER 1981, fig. 5), as well as further occurrences of the latter species and *B. taichoutense* KRÖGER & LEFEBVRE, from horizons regarded as of early to mid-Floian age in Morocco (KRÖGER & LEFEBVRE 2012) make for difficulty in identifying the ancestor of *Bathmocerases*. Nevertheless, when all records are considered, with the exception of South

America, *Bathmocerases* is present in western Gondwana from the mid-Floian to the Darriwilian and is also present in the Southern China and the Baltic during the Dapinginian and Darriwilian.

Margaritoceras CECIONI & FLOWER from the Floian of Bolivia and *Sarcodosoceras* EVANS from the Darriwilian of the Welsh Basin are similar, although the latter differs in its lower rate of conch expansion, narrower siphuncle, and compressed ovoid cross-section (EVANS 2005). Although they are likely to be closely related, *Margaritoceras* is at present known only from the Central Andean Basin of Bolivia and northwest Argentina, where *M. diploide* is now known to be present in the mid Floian possibly late Floian (CICHOWLSKI unpublished data) whilst a further species of *Margaritoceras* is known from the late Floian (CICHOWLSKI unpublished data). Additional, undescribed taxa are known from the mid-Floian of the Central Andean Basin (CICHOWLSKI unpublished data). As it stands, the evidence from South America suggests that the Eothinoceratidae underwent a radiation at least during the Floian that appears to be restricted to South America, although taxa such as *Sarcodosoceras* may suggest that some of these lineages later extended into Avalonia.

		Balto-scandian Stages	Swedish Formations	Trilobite zonation	Lituitid zonation	Graptolite zonation	Conodont zonation
Darrivillian	4c	Uhakuan (part)	Furudal Limestone	<i>Iliaenus crassicaudata</i>	<i>Ancistroceras undulatum</i>	<i>Hustedograptus teretiusculus</i>	<i>Pygodus anserinus</i>
		Lasnamägian	Folkeslunda Limestone	<i>Iliaenus schroeteri</i>	<i>Lituites perfectus</i>	<i>Pseudoamplexograptus distichus</i>	<i>Pygodus serra</i>
			Seby Limestone		<i>Lituites lituus</i>		
			Skärlov Limestone		<i>Trilacnocras discors</i>		
		Aserian	Segerstad Limestone	<i>Iliaenus planifrons/ Iliaenus platyurus</i>	<i>'Lituites' toernquisti</i>	<i>Pterograptus elegans</i>	<i>Eoplacognathus suecicus</i>
					<i>Angelinocras latum</i>		
	4b	Kundan	Holen Limestone	<i>Megistaspis gigas</i> <i>Megistaspis obtusicauda</i>		<i>Nicholsongraptus fasciculata</i>	<i>Eoplacognathus pseudoplanus</i>
	4a			<i>Asaphus raniceps</i>	<i>Holmiceras praecurrens</i>	<i>Holmograptus lentus</i>	<i>Yangtzeplacognathus crassus</i>
				<i>Asaphus expansus</i>		<i>Undulograptus austrodentatus</i>	<i>Lenodus variabilis</i>

Fig. 3: Stratigraphical distribution of lituitid species in Baltica during the Darrivillian, demonstrating the high resolution provided by these taxa in comparison with trilobites, graptolites and conodonts.

The Eothinoceratidae and Bathmoceratidae were widely distributed around Gondwana during the Early and Middle Ordovician, and where the range of a species is known, may resolve to a stage or even a timeslice (Fig. 2). As some inshore assemblages of high latitude Gondwana (e.g. parts of the early Floian, Bolahaul Member of the Ogof Hên Formation of Wales) may be dominated by molluscs, whilst graptolites and trilobites are relatively rare (e.g. FANG & COPE 2004), these cephalopods may have the potential to constrain ages in some cases.

3. Lituitida. Lituitid nautiloids are common and distinctive components of Middle Ordovician cephalopod faunas in the 'Orthoceratite Limestone' facies of Baltoscandia (especially Sweden) and coeval carbonate sequences, such as the Dawan Formation in China. They are virtually cosmopolitan in occurrence, and are known from the USA and Newfoundland (FLOWER 1975), Wales (EVANS 2005, in prep.), China (e.g.

YU 1930; LAI 1982; LAI 1986; QI 1980), Korea (YUN 1999, 2002), Estonia (BALASHOV 1953), Norway (SWEET 1958), Sweden (ANGELIN & LINDSTRÖM 1880; HOLM 1891; KING 1999) and the 'Diluvium-Geschiebe' of northern Poland (DZIK 1984) and Germany (BOLL 1857; NOETLING 1884; REMELÉ 1880, 1881). A substantial number of lituitid genera have been described since the 19th century (e.g. HYATT 1894; TEICHERT et al. 1964) and the taxa are firmly established within cephalopod literature.

The Swedish lituitid fauna extends mainly from the early Kundan to Uhakuan stages. This general succession of taxa closely corresponds with lituitid faunas described elsewhere, especially China (CHEN & LIU 1976). In the Swedish 'Orthoceratite Limestone' sequence, biostratigraphical data from the latest Volkhovian and earliest Kundan stages of Öland and Östergötland indicate that the weakly cyrtconic Sinoceratidae occur below the coiled Lituitidae. The former are represented by




Fig. 4: Occurrence ranges of *Cenoceras* species in the Early Jurassic of southern England, compared with the standard ammonite zonation. Inset: *Cenoceras intermedium* (SOWERBY, 1816), reproduced from D'ORBIGNY (1843: pl. 27, fig. 1).

Rhynchorthoceras and related genera that extend up into the Aserian and Lasnamagian stages. The earliest Litu-
itidae occur in the mid Kundan and are represented by
forms such as *Holmiceras praecurrens* (HOLM, 1891). *An-*
cistroceras, with its very rapidly expanding conch is
recorded from the late Kundan of Kinnekulle,
Västergötland, and extends through the Aserian and
Lasnamagian, but is more typical of and numerous in
the late Lasnamagian and Uhakuan stages. *An-*
gulinoceras latum (ANGELIN & LINDSTRÖM, 1880) is
recorded only from the lower part of the Segerstad
Limestone (Aserian) and the maximum development
and diversity of Litu-
itidae is attained within the Las-
namagian-aged Seby and Folkeslunda limestones where
various species of *Litu-
ites*, *Trilacino-*
ceras and *Cyclolitu-*
ites are of common occurrence.

Apart from the possession of complex sutures, liti-
itids possess all the attributes typically associated with
ammonoids (especially ammonites) which make them
such valuable and reliable biozonal indicators and capa-
ble of biostratigraphic correlation with a high degree of
resolution. Namely, lituitids possess distinctive, easily
recognisable conchs whose form and ornament enables
taxa to be readily distinguished; they evolved rapidly
and have a wide geographical occurrence; and although
they are more commonly found in shallow water, plat-
form carbonate sequences, they also occur in mudstone
facies, such as those of the Llanfawr Mudstone Forma-
tion of central Wales (EVANS in prep).

Within Sweden, the use of some lituitid taxa as
stratigraphical indicators has long been recognised (e.g.
JAANUSSON & MUTVEI 1953; JAANUSSON 1963) and
their common occurrence in some beds has given rise to
former names such as 'Ancistroceras Limestone' (now
termed Furudal Limestone). Closer examination of the
Darriwilian lituitid faunas (KING 1990, unpublished and
in prep.) demonstrates that these lituitids have consid-
erable potential as zonal fossils, and achieve a resolution
at least as good as, and for the Aserian and Lasnamagian
portion of the succession, potentially finer than that
currently recognised using established trilobite, graptolite
or conodont zonal schemes (Fig. 3).

Silurian-Devonian

In recent years concentrated efforts have been made
to improve the knowledge of the distribution and tax-
onomy of Silurian nautiloid cephalopod faunas and
many existing collections have been revised using up to
date taxonomic criteria as well as collection of new ma-
terial from horizons with precise biostratigraphic data.
Consequently systematic studies of Silurian nautiloid
cephalopods from a variety of geographical settings and
the observed temporal and spatial data from these fau-
nas may now be considered a reliable tool for palaeobio-
geographical reconstruction. In Europe the main work
on Silurian faunas has been done in the British Isles
(EVANS 1994; EVANS & HOLLAND 1995; HOLLAND 1998,
1999, 2000a-c, 2002, 2003, 2004, 2007, 2010; HOL-

LAND & STRIDSBERG 2004), Sweden (STRIDSBERG 1985), Prague Basin (DZIK 1984; GNOLI 1997; KOLEBABA 1975, 1977, 1999, 2002; MANDA 1996, 2008; MANDA & KŘÍŽ 2006, 2007; MANDA & TUREK 2009a,b; MAREK 1971; MAREK & TUREK 1986; STRIDSBERG & TUREK 1997; TUREK 1975, 2008), South West Sardinia (GNOLI 1990; GNOLI & SERPAGLI 1977, 1991; GNOLI & SERVENTI 2006, 2009 and references therein; SERPAGLI & GNOLI 1977), Spain (BOGOLEPOVA 1998a), France (RISTEDT 1968; SERVENTI & FEIST 2009) the Carnic Alps of Austria and Italy (BOGOLEPOVA 1998b; GNOLI & HISTON 1998; GNOLI & SERVENTI 2008; GNOLI et al. 2000; HISTON 1997, 1998, 1999a, b, 2002, 2012a, b; RISTEDT 1968, 1969, 1971; SERVENTI & GNOLI 2000; SERVENTI et al. 2006, 2010) and the Graz Palaeozoic of Austria (HISTON et al. 2010). Tentative correlations are now possible between Avalonia (British faunas), some areas positioned along the Northern Gondwana Margin (Carnic Alps, Sardinia, France and Spain) and Bohemia and Baltica, although problems do still exist in recognition of faunas at both generic and specific level due to poor preservation and lack of precise taxonomic diagnoses.

Detailed study of Silurian-Devonian nautiloid faunas from Morocco by (KRÖGER 2008) presented together with precise stratigraphic and lithofacies data for the collection localities has highlighted further exchange of faunas between Peri-Gondwana Terranes. Studies of particular taxa and nautiloid biodiversity, again within precise stratigraphic biozones and detailed facies studies, in the Silurian of the Prague Basin in relation to palaeobiogeographical distribution and oceanic states of the North Gondwana area and Perunica by MANDA et al. (2009, 2010) and others (see list above) are important contributions to these fields and confirms that nautiloid cephalopods are indeed reliable indicators for this interval. Studies of faunas at both a local and regional scale within single biozones are still preferable for building a consistent database for future reference.

Certain taxa show potential as biostratigraphical indicators for the Silurian-Devonian interval as has been shown by various studies (see list above): *Orthocycloceras*, *Hemicosmorthoceras*, *Plagiostomoceras*, *Pseudocycloceras*, *Columenoceras*, *Parakionoceras*, *Kionoceras*, *Dawsonoceras*, *Sphaerorthoceras*, *Temperoceras* as well as representatives of the Phragmocerids, Oncocerids and Tarphycerids such as *Ophioceras*. Nautiloid cephalopod assemblages were broadly defined by GNOLI & SERPAGLI (1991) and by MANDA & KŘÍŽ (2006) using a suite of taxa which dominated in certain Silurian series: *Pseudocycloceras transiens-Columenoceras grande* (Wenlock–early Ludlow), *Merocycloceras declive-Cryptocycloceras deludens* (early Ludlow), *Kopaninoceras thyrus-Orthocycloceras fluminese* (late Ludlow–Pridoli/Early Devonian, *Pseudocycloceras duponti-*

Sphooceras truncatum (Wenlock), *Pseudocycloceras duponti-Kionoceras doricum* (early Ludlow) and *Pseudocycloceras nereidum-Sphooceras truncatum* (Ludlow). Taxa such as *Deiroceras* and *Jovellania* have been seen to dominate in certain intervals of the Devonian in Morocco (KRÖGER 2008) and may prove useful for recognition of marker beds deposited within sequences.

There have been numerous studies of the well-known Silurian-Devonian in age ‘Orthoceras limestones’ or Cephalopod Limestone Biofacies both with regard to their depositional cycles and biotic content. The Carnic Alps of Austria is a key locality along the Northern Gondwana margin regarding Silurian biostratigraphical correlation where the Silurian Cephalopod Limestone Biofacies is well preserved. The Cellon section has been utilized as a geographic reference district (RD) for both conodont correlation studies (KLEFFNER 1989, 1995) and for evaluation of global eustatic changes (JOHNSON 2010) for the North Gondwana area. Recognition of environmental and water depth changes based on the fossil assemblages (mostly trilobites, brachiopods and bivalves) from the Silurian depositional sequences developed there (BRETT et al. 2009) places a tight control on small scale bioevents within well-defined conodont (WALLISER 1964), graptolite (JAEGER 1975) and chitinozoan (PRIEWALDER 1997) biozones. Particular emphasis has been placed in these studies on establishing the response of marine faunas to oscillations in sea-level and to the oceanic variations (chemistry, temperature, currents) recorded (WENZEL 1997; KŘÍŽ 1998, KŘÍŽ et al. 2003) during this time interval on a local scale for comparison with data from other North Gondwana terranes such as Sardinia and Bohemia and on a global scale with some sectors of Avalonia (the British Isles) and Laurentia (North America).

Correlation of the nautiloid faunal assemblages from the cephalopod limestone biofacies levels and their taphonomic signatures within the contexts outlined above with evidence for pronounced redox changes, surface currents, regression/transgression sequences within precise intervals from the Carnic Alps (Austria) successions may identify common controlling factors in the palaeogeographic distribution and migrational routes of these faunas (HISTON 2012b). Current studies, in line with those of MANDA (2009), MANDA & FRYDA (2010) and others, attempt to identify controlling factors on a local scale for nautiloid distribution within precise time slices that may then be recognized in other areas where these faunas occur along the North Gondwana Margin (HISTON 2012a). This is an on-going study done in parallel with revision of historical collections (GNOLI & HISTON 1998; HISTON 1999; GNOLI et al. 2000), systematic collection and description within precise biozones

(HISTON 2002; HISTON et al. 2010) and taphonomic studies (HISTON 1999, 2002, 2012a and references therein). This holostratigraphical approach may provide further evidence for the reconstruction of reliable nautiloid assemblages or identification of precise marker taxa.

Carboniferous

The monographic works by HYATT, MILLER and others in the USA, SHIMANSKY in Russia, FOORD and TURNER in the British Isles, and DE KONINCK from Belgium constituted the foundations for the study on Carboniferous nautiloid cephalopods (see papers listed under references). In recent years some taxonomic revision of these nautiloids has been carried out (see studies listed by HISTON, MAPES, NIKO, STURGEON and others), but the use of rapidly evolving goniatite lineages to establish a biostratigraphy, particularly for the Upper Carboniferous is predominant and consideration of the stratigraphical potential of nautiloid cephalopods has been neglected. However, several taxa are worthy of mention and are known to be abundant at certain intervals in a diverse range of palaeogeographical settings. Large specimens of *Rayonnoceras* are known to be markers within Lower Carboniferous strata of Europe and the USA where they are common in the Brigantian/Mississippian. The annulated *Cycloceras* and taxa of the Oncocerida such as *Poterioceras* and *Welleroceras* are common in the Lower Carboniferous. Several Pseudorthocerataceae taxa are restricted to the Carboniferous such as *Mitorthoceras*. The distinctively ornamented *Brachycycloceras* is a common element in the Upper Carboniferous. A variety of coiled nautiloid taxa such as *Vestinautilus*, *Trigonoceras*, *Aphelaeceras*, *Epistroboceras*, *Maccocyoceras*, *Asymptoceras*, *Acanthonautilus* and *Bistrialites* are typical of the Lower Carboniferous of the British Isles, Belgium, Asia and the USA. More precise biostratigraphical data may be obtained through further study of the ranges of these taxa at species level. However, to date, such a zonal scheme has not been developed.

Permian

The papers by MILLER et al. (1942, 1949) of the Permian nautiloids of the USA still stand as reference works for taxonomy of the group. However, there are few studies in relation to the development of the biostratigraphical potential of nautiloid cephalopods for this interval. An investigation of the Late Permian nautiloid faunas of the Bellerophon Formation from the Dolomites of Northern Italy (POSENATO & PRINOTH 2004, 2007; POSENATO 2010) indicates that certain nautiloid taxa representing species of *Tainoceras*, *Tirolonauutilus*, *Liroceras* and *Foordiceras* may indeed be used as markers for precise levels within the succession whilst

some species may be assigned lineages and may be compared with assemblages described by TEICHERT & KUMMEL (1973) from the Late Permian of northwest Iran.

Triassic

Despite the relative paucity of works on Triassic nautiloid cephalopods, there is strong evidence of their potential value to some aspects of Triassic biostratigraphy. KUMMEL (1953) reviewed many Triassic nautiloid genera, indicating their stratigraphical ranges and a potential for their biostratigraphical use. This potential has been realised through the development of a detailed zonal scheme using nautiloids (SOBOLEV 1994) for the Boreal Triassic of Siberia. Such a scheme clearly demonstrates the value of the organisms as biostratigraphical tools.

Jurassic and Cretaceous

After the disappearance of seven families and approximately thirty genera during the Late Triassic extinctions (KUMMEL 1964, fig. 294), only the Nautilidae, represented by the single genus *Cenoceras* is conventionally regarded as having survived into the Jurassic where it underwent a radiation that led to all subsequent nautiloid genera including *Nautilus*. Analysis of the distribution of Middle Jurassic nautilids in western France by BRANGER (2004) indicates that the ranges of some taxa may be no more than one or two ammonite zones, indicating a potential utility as biostratigraphical tools. The same appears to be true for Lower Jurassic nautilids.

Historically the name *Cenoceras* has been broadly applied to a diverse range of Late Triassic to early Middle Jurassic nautilids that constitute a “plastic evolving complex” (KUMMEL 1956, p. 361). It is now recognised that this masks a number of distinct lineages within a rapidly evolving nexus of Lower Jurassic nautilid faunas (TINTANT 1984, 1987; RULLEAU 2008). Previous workers (HYATT 1894; SPATH 1927) recognised a number of different morphotypes within the ‘*Cenoceras* complex’ and provided names for several forms (including *Digonioceras*, *Ophionauutilus* and *Sphaeronautilus*). The taxonomic status of these taxa requires revision.

More recently, TINTANT (1984) described three subgenera within *Cenoceras*: *Cenoceras* (in a restricted sense) for forms with relatively stout, involute conchs possessing a spiral ornament on the ventral and lateral surfaces of the whorls; *Hemicenoceras* for compressed forms possessing spiral ornament that is mainly confined to the ventral area, and *Metacenoceras* for forms with a flattened venter and an ornament consisting only of weak transverse growth lines. The earliest *Metacenoceras* occur in the early Sinemurian (Shales-with-Beef

Member, Charmouth Mudstone Formation near Lyme Regis, Dorset, UK [KING unpublished data]); *Hemicenoceras* is typically first encountered in the Carixian, where it is represented by *H. arare* (DUMORTIER, 1869) and *H. egregium* (PIA, 1914). In this paper, *Cenoceras* is used in the restricted sense employed by TINTANT.

The Hettangian to Carixian sequence of *Cenoceras malherbii* (TERQUEM, 1855) – *C. intermedium* (SOWERBY, 1816) – *C. striatum* (SOWERBY, 1817) – *C. pertextum* (DUMORTIER, 1867) – *C. simillimum* (FOORD & CRICK, 1890) was recognised in French successions by TINTANT (1984). In the United Kingdom, the same sequence of taxa is present (in part) in the Lower Jurassic of South Wales, and on the North Somerset and Dorset coasts (Fig. 4). Within this lineage there is a tendency for conchs to become more involute (from 20% in Hettangian forms to <10% in Carixian taxa); for the siphuncle to migrate from a ventrocentral to central position; for the whorl section to become more rounded and arched, and less quadrate; and for suture lines to become more sinuous, culminating with *C. jourdani* (DUMORTIER, 1874) in the Toarcian (KING 2011).

Late Hettangian and early Sinemurian strata on the Somerset coast (particularly within the *angulata* and *bucklandi* zones) include several monotonous beds of dark grey to black shales and bituminous mudstones which are virtually devoid of ammonites. These units contain fossil nautilids that are embedded in the substrate at orientations that range from horizontal to vertical, and provided hard attachment surfaces for a range of organisms of which oysters and crinoids are the most abundant.

Wherever ammonites are present in the Lower Jurassic, by comparison, nautilids make relatively poor biostratigraphical indicators. However, when ammonites are absent, cenoceratid nautilids can be used for biostratigraphical purposes and provide a resolution of at least stage or even substage level. Unlike Middle Jurassic nautilids (BRANGER 2004), there are, at present, no known Lower Jurassic cenoceratid taxa that are characteristic indicators of an individual ammonite zone, although further research may change this picture.

Cretaceous nautilids from southern South America are relatively scarce and of low diversity. Most belong to one or two genera: *Cymatoceras* HYATT, and *Eutrephoceras* HYATT. Both genera possess a strongly involute, inflated and globose conch with an orthochoanitic, sub-central siphuncle, as well as simple sutures. Both genera are cosmopolitan in distribution, but at species level, cosmopolitan distributions are almost nonexistent (CICHOWOLSKI 2003) which may reflect the nektobenthic habit of these animals, as well as the lack of a plankton-

ic larval stage. Furthermore, the skeletal morphology of nautilids in general, and of these genera in particular, is markedly homogeneous with respect to the characters used to diagnose different species.

All these considerations suggest that attempts to use Cretaceous nautilids for biostratigraphical purposes are likely to be disappointing. This is compounded when the long stratigraphical ranges of some of the species are taken into consideration. *Cymatoceras perstriatum* (STEUER) is a typical example of such a species. *C. perstriatum* (STEUER) occurs in strata of Tithonian-Hauterivian age in the Neuquén Basin (CICHOWOLSKI 2003); the late Tithonian-early Berriasian of the Chilean Aconcagua platform (CORVALÁN 1959; BIRÓ-BAGÓCZKY 1964), and the late Valanginian-early Albian of the Chañarcillo Basin (northern Chile: HOFSTETTER et al. 1957; SEGERSTROM 1960). The whorl cross-section, the sutural pattern, and overall shape of the conch all exhibit a marked ontogenetic and intraspecific variability (CICHOWOLSKI 2003). Such a range of variation implies that an extra effort may be required in order to positively identify individual members of this particular taxon.

However, there are some nautilid species possessing easily recognisable morphologies that present low variability within and between individuals. When such taxa have a relatively short stratigraphical range, and other biostratigraphic markers are lacking, they may provide valuable information with regard to the age of the strata in which they are found. One such example may be provided by *Eutrephoceras dorbygnianum* (FORBES in DARWIN, 1846), known from the Antarctic Peninsula, Quiriquina Island in Chile, the Austral Basin of southern Argentina, and possibly from Angola (STEINMAN 1895; SPATH 1953; HOWARTH 1965; STINNESBECK 1986; CICHOWOLSKI et al. 2005; NIELSEN & SALAZAR 2011). This species is characterized by an almost straight suture, combined with a small, acute, umbilical saddle, and a semilunate whorl cross-section (CICHOWOLSKI et al. 2005). Based on its occurrence, this species ranges through the Campanian and Maastrichtian, becoming particularly abundant during the Maastrichtian. Thus, like the Lower Jurassic *Cenoceras*, the stratigraphical range of some Late Cretaceous nautilids may resolve to one or two stages.

Concluding remarks

This survey of the use and potential use of nautiloid cephalopods in biostratigraphy is inevitably, incomplete. Nevertheless, examples provided by the liutitids (above) and the Triassic nautiloids of Siberia (SOBOLEV 1994) demonstrate that when diversification rates are high, these organisms may be used to define biostrati-

graphical schemes to resolutions that are at least equivalent to those generated through the use of other groups of organisms. In situations where other groups of organisms are rare or not present, nautiloid cephalopods may provide an alternative basis for a biostratigraphical scheme. Finally, and perhaps more significantly, nautiloid cephalopods are powerful palaeobiogeographical tools, especially, but not exclusively during the Ordovician and Silurian. In order to refine their value as palaeobiogeographical tools, it will be necessary also to further refine their biostratigraphy. This is a task that will continue through the study of well-documented collections as well as newly collected material that may help resolve questions related to older collections, and bring to light new material from poorly known areas.

References

- ADRAIN J.M. (2011): A new trilobite stratigraphy for the Lower Ordovician of western Laurentia and prospects for international correlation using pelagic trilobites. — In: GUTIÉRREZ-MARCO J.C., RÁBANO I. & D. GARCÍA-BELLIDO (Eds), *Ordovician of the World*. Cuad. Mus. Geomin. **14**: 41-42.
- ANGELIN N.P. & G. LINDSTRÖM (1880): *Fragmenta Silurica e dono Caroli Henrici WEGELIN*. — Samson & Wallin, Stockholm: 60 pp.
- BALASHOV Z.G. (1953): Coiled and semicoiled Nautiloidea of the Ordovician of the Baltic area. — Trudy Vsesoyuznogo Nauchno-Issled Geol.-Razvedo Inst. (VNIGRI) **78**: 217-268. [In Russian]
- BARRANDE J. (1865-1877): *Système Silurien du centre de la Bohême*. Part 2. Céphalopodes. — Prague: part 1, p. 1-712, 1867; part 2, p. 1-263, 1870; part 3, p. 1-804, 1874; part 4, p. 1-743, 1877; part 5, p. 1-743, 1877; supplement p. 1-297, 1877; pls 1-107, 1865; pls 108-244, 1866; pls 245-350, 1868; pls 351-460, 1868; pls 351-460, 1870; supplement pls 461-544, 1877.
- BECKER R.T. (2000): Taxonomy, evolutionary History and Distribution of the middle to late Famennian Wocklumeriina (Ammonoidea, Clymeniida). — Fossil Rec. **3**: 27-75.
- BIRÓ-BAGÓCZKY L. (1964): Estudio sobre el límite entre el Titoniano y el Neocomiano en la Formación Lo Valdés, Provincia de Santiago, principalmente en base a ammonioideos. — Universidad de Chile, Departamento de Geología, memoria de título: 118 p.
- BLAKE J.F. (1882): *A Monograph of the British Fossil Cephalopoda*. Pt. 1. Introduction and Silurian Species. — J. Van Voorst, London: 248 pp.
- BOGOLEPOVA O.K. (1998a): A brief account on the Upper Silurian cephalopods from the Valle syncline, province of Seville (Ossa Morena Zone, southern Spain). — In: GUTIÉRREZ-MARCO J.C. & I. RABANO (Eds), *Proceedings 6th International Graptolite Conference & 1998 Field Meeting, IUGS Subcommission on Silurian Stratigraphy*. Temas Geol.-Min. ITGE **23**: 63-66.
- BOGOLEPOVA O.K. (1998b): Silurian cephalopods: New data from the Carnic Alps of Austria. — In: GUTIÉRREZ-MARCO J.C. & I. RABANO (Eds), *Proceedings 6th International Graptolite Conference & 1998 Field Meeting, IUGS Subcommission on Silurian Stratigraphy*. Temas Geol.-Min. ITGE **23**: 60-62.
- BOLL E. (1857): Beitrag zur Kenntnis der silurischen Cephalopoden im norddeutschen Diluvium und den anstehenden Lager Schwedens. — Arch. Ver. Freunde d. Naturgesch. Mecklenburg **11**: 58-95.
- BOSTON W.B. & R.H. MAPES (1991): Ectocochliate cephalopod taphonomy. — In: DONOVAN S.K. (Ed.), *The Processes of Fossilization*. Belhaven Press, London: 220-240.
- BRANGER, P. (2004): Middle Jurassic Nautiloidea from Western France. — Riv. Italiana Paleont. Stratigr. **110**: 141-149.
- BRETT C.E., FERRETTI A., HISTON K. & H.P. SCHÖNLAUB (2009): Silurian sequence stratigraphy of the Carnic Alps, Austria. — Palaeogeogr., Palaeoclimatol., Palaeoecol. **279**: 1-28.
- BUCKMAN S.S. (1898): On the grouping of some divisions of so-called „Jurassic“ time. — Quart. J. Geol. Soc. **54**: 442-462.
- CALLOMON J.H., DIETL G. & K.N. PAGE (1989): On the Ammonite faunal horizons and Standard Zonation of the Lower Callovian Stage in Europe. — In: RACHA R.B. & A.F. SOARES (Eds), *2nd International Symposium on Jurassic Stratigraphy*. Centro de Estratigrafia Paleobiologia de Universidade Nova de Lisbon, Lisbon: 17-18.
- CATALANI J.A. (1987): Biostratigraphy of the Middle and Late Ordovician Cephalopods of the Upper Mississippi Valley area. — In: SLOANE R.E. (Ed.), *Middle and Late Ordovician Lithostratigraphy and Biostratigraphy of the Upper Mississippi Valley*. Minnesota Geol. Survey, Rep. Investig. **35**: 187-189.
- CECIONI G. & R.H. FLOWER (1985): Bathmoceratidae (Nautiloidea, Ordovicio) de Sud América. — N. Jahrb. Geol. Paläont. Abh. **170**: 343-357.
- CHEN J.-Y. & G.W. LIU (1976): Ordovician nautiloids. — Mem. Nanjing Inst. Geol. Palaeont. **7**: 42-53.
- CHEN J.-Y. & C. TEICHERT (1987): The Ordovician cephalopod Suborder Cyrtocerina (Order Ellesmerocerida). — Palaeont. Cathayana **3**: 145-229.
- CICHOWOLSKI M. (2003): The nautiloid genus *Cymatoceras* from the Cretaceous of the Neuquén and Austral basins, Argentina. — Cretaceous Res. **24**: 375-390.
- CICHOWOLSKI M. (2009): A review of the endocerid cephalopod *Protocyrtendoceras* from the Floian (Lower Ordovician) of the Eastern Cordillera, Argentina. — Acta Palaeont. Polonica **54**: 99-109. doi:10.4202/app.2009.0110
- CICHOWOLSKI M. & N.E. VACCARI (2011): The oldest record of Eothinoceratidae (Ellesmerocerida, Nautiloidea): Middle Tremadocian of the Cordillera Oriental, NW Argentina. — Geol. J. **46**: 42-51.
- CICHOWOLSKI M., AMBROSIO A. & A. CONCHEYRO (2005): Nautilids from the Upper Cretaceous of the James Ross Basin, Antarctic Peninsula. — Antarctic Sci. **17** (2): 267-280. doi:10.1017/S0954102005002671
- COCKS L.R.M. & T.H. TORSVIK (2002): Earth geography from 500 to 400 million years ago: a faunal and palaeomagnetic review. — J. Geol. Soc. London **159**: 631-644.
- COOPER B.J. (1981): Early Ordovician conodonts from the Horn Valley Siltstone, central Australia. — Palaeontology **24**: 147-183.
- CORVALÁN J. (1959): El Tithoniano de Río Leñas, Provincia de O'Higgins, con una revisión del Tithoniano y Neocomiano de la parte Chilena del geosinclinal andino. — Bol. Inst. Invest. Geol. **3**: 1-63.
- CRICK R.E. (1980): Integration of paleobiogeography and paleogeography: evidence from Arenigian nautiloid biogeography. — J. Paleont. **54**: 1218-1236.

- CRICK R.E. (1988): Buoyancy regulation and macroevolution in nautiloid cephalopods. — *Senckenberg. lethaea* **69**: 13-42.
- CRICK R.E. (1990): Cambro-Devonian biogeography of nautiloid cephalopods. — In: MCKERROW W.S. & C.R. SCOTSE (Eds), *Palaeozoic Palaeogeography and Biogeography*. Geol. Soc. Mem. **12**: 147-161.
- CRICK R.E. (1993): Biogeography of early and middle Paleozoic nautiloid cephalopods: evidence for barriers to dispersal and evolution. — *Geobios* **15**: 91-105.
- DARWIN C.R. (1846): Geological observations on South America. Being the third part of the geology of the voyage of the Beagle, under the command of Capt. FITZROY, R.N. during the years 1832 to 1836. — Smith Elder & Co., London: 280 pp.
- DE KONINCK L.G. (1842-44): Description des animaux fossiles qui se trouvent dans le terrain Carbonifere de la Belgique. — Liege: 650 pp.
- DE KONINCK L.G. (1878-1882): Faune du calcaire carbonifere de la Belgique. — Musee Royal d'Histoire Naturelle de Belgique, Bruxelles: pt. 1 (1878): 152 pp., pt. 2 (1880): 133 pp.
- DUMORTIER E. (1869): Etudes paléontologique les sur les dépôts jurassiques du bassin du Rhone. Pt. 3. Lias Moyen. — Savy Ed., Paris: 348 pp.
- DUMORTIER E. (1874): Etudes paléontologique les sur les dépôts jurassiques du bassin du Rhone. Pt. 4. Lias Moyen. — Savy Ed., Paris: 339 pp.
- DZIK J. (1984): Phylogeny of the Nautiloidea. — *Palaeont. Polonica* **45**: 1-219.
- EVANS D.H. (1994): Irish Silurian Cephalopods. — *Irish J. Earth Sci.* **13**: 113-48.
- EVANS D.H. (2005): The Lower and Middle Ordovician cephalopod faunas of England and Wales. — *Monogr. Palaeontogr. Soc.*, London **158**: 1-81.
- EVANS D.H. (2007): A Middle Ordovician cephalopod fauna from Cuzco Province, southern Perú and its palaeobiogeographical significance. — *Geol. J.* **42**: 25-36.
- EVANS D.H. (2011): The Lower Ordovician cephalopod faunas of the Durness Group, north-west Scotland. *Monogr. Palaeontogr. Soc.* London **165**: 1-131.
- EVANS D.H. & C.H. HOLLAND (1995): The nautiloid cephalopod order Endocerida in the Silurian. — *Paläont. Z.* **69**: 343-352.
- FANG Z.-J. & J.W.C. COPE (2004): Early Ordovician bivalves from Dali, West Yunnan, China. — *Palaeontology* **47**: 1121-1158.
- FLOWER R.H. (1946): Ordovician cephalopods of the Cincinnati region. Part I. — *Bull. American Paleont.* **29**: 1-751.
- FLOWER R.H. (1964): The nautiloid order Ellesmeroceratida (Cephalopoda). — *New Mexico Inst. Min. Technol. Mem.* **12**: 1-234.
- FLOWER R.H. (1975): American Lituitidae (Cephalopoda). — *Bull. American Paleont.* **67**: 139-173.
- FLOWER R.H. (1976): Ordovician cephalopod and their role in correlation. — In: BASSETT M.G. (Ed.), *The Ordovician System: Proceedings of a Palaeontological Association Symposium*, Birmingham, September 1974. University of Wales Press and National Museum of Wales, Cardiff: 523-555.
- FLOWER R.H. (1985): Progress and changing concepts in cephalopod and particularly nautiloid phylogeny and distribution. — In: WEIDMANN J. & J. KULLMANN (Eds), *Cephalopods: Present and Past*. 2nd International Cephalopod Symposium. Stuttgart. Schweizerbart'sche Verlagsbuchhandlung, Stuttgart: 18-24.
- FOERSTE A.F. (1932): Black River and other cephalopods from Minnesota, Wisconsin, Michigan and Ontario. Pt. 1. — *Denison Univ. Bull., J. Sci. Lab.* **27**: 47-136.
- FOERSTE A.F. (1933): Black River and other cephalopods from Minnesota, Wisconsin, Michigan and Ontario. Pt. 2. — *Denison Univ. Bull., J. Sci. Lab.* **28**: 1-164.
- FOORD A.H. (1888-1891): Catalogue of the Fossil Cephalopoda in the British Museum (Natural History) London. — *British Museum (Natural History)*, London: Pt. 1, 344 pp.; Pt. 2, 407 pp.
- FOORD A.H. (1897-1903): Monograph of the Carboniferous Cephalopods of Ireland. — *Palaeontographical Society*, London: 1-234.
- FOORD A.H. & G.C. CRICK (1890): Descriptions of new and imperfectly-defined species of Jurassic Nautili contained in the British Museum (Natural History). — *J. Nat. Hist., ser. 6* **5**: 265-291.
- FOORD A.H. & G.C. CRICK (1897): Catalogue of the Fossil Cephalopoda in the British Museum (Natural History) London. Pt. 3. — *British Museum (Natural History)* London: 303 pp.
- FREY R.C. (1995): Middle and Upper Ordovician nautiloid cephalopods of the Cincinnati arch region of Kentucky, Indiana, and Ohio. — *U.S. Geol. Survey Prof. Paper* **1066-P**: 1-126.
- FURNISH W.M. & B.F. GLENISTER (1964): Nautiloidea-Tarphycerida. — In: MOORE R.C. (Ed.), *Treatise on Invertebrate Paleontology*. Part K. Mollusca 3, Cephalopoda. General features, Endoceratoidea, Actinoceratoidea, Nautiloidea, Bactritoidea. Geological Society of America and University of Kansas Press, Boulder, Colorado and Lawrence: 343-368.
- GNOLI M. (1990): New evidence of faunal links between Sardinia and Bohemia in Silurian time on the basis of nautiloids. — *Boll. Soc. Paleont. Italiana* **29** (3): 289-307.
- GNOLI M. & K. HISTON (1998): Silurian nautiloids from the Carnic Alps: a preliminary investigation. — *Boll. Soc. Paleont. Italiana* **36**: 311-330.
- GNOLI M. & E. SERPAGLI (1977): Silurian cephalopods of the MENEGHINI collection (1857) – with reproduction of the original plate. — *Boll. Soc. Paleont. Italiana* **16** (2): 137-142.
- GNOLI M. & P. SERVENTI (2006): A further oncocerid nautiloid from the upper Silurian of southwest Sardinia. — *Geo.Alp* **3**: 3-8.
- GNOLI M. & P. SERVENTI (2008): A new cephalopod from the early Silurian of the Carnic Alps (Italian side). — *Riv. Italiana Paleont. Stratigr.* **114** (2): 171-178.
- GNOLI M. & P. SERVENTI (2009): Silurian nautiloid cephalopods from Sardinia: the state of the art. — In: CORRADINI C., FERRETTI A. & P. STORCH (Eds), *The Silurian of Sardinia*. *Rendiconti Soc. Paleont. Italiana* **3** (1): 109-118.
- GNOLI M., HISTON K. & P. SERVENTI (2000): Revision of the Silurian Cephalopods from the Carnic Alps: The Gortani and Vinasca de Regny collection, 1909. — *Boll. Soc. Paleont. Italiana* **39** (1): 3-12.
- HALL J. (1847): *Palaeontology of New York*. Vol. I. Descriptions of the Organic Remains of the Lower Division of the New York System. — C. Van Benthuysen, Albany: 338 pp.
- HERWIG P. & R. POSENATO (2007): Late Permian Nautiloids from the Bellerophon Formation of the Dolomites (Italy). — *Palaeontographica A* **282** (1-6): 135-165.

- HEWITT R.A. & G.E.G. WESTERMANN (1996): Post-mortem behaviour of early Paleozoic nautiloids and paleobathymetry. — *Paläont. Z.* **70**: 405-424.
- HISTON K. (1997): Cephalopod limestones. Cellon Section. — *Ber. Geol. Bundesanst.* **40**: 92-99.
- HISTON K. (1998): Die Nautiloideen-Fauna aus dem Silur der Karnischen Alpen. — *Geol.-paläont. Mitt. Innsbruck* **23**: 105-115.
- HISTON K. (1999a): Revision of Silurian nautiloid cephalopods from the Carnic Alps (Austria) – The HERITSCH (1929) Collection in the Geological Survey of Austria. — *Abh. Geol. Bundesanst.* **56** (1): 229-258.
- HISTON K. (1999b): Silurian cephalopod limestone facies in the Carnic Alps (Rauchkofel Boden section, Austria): taphonomy of the nautiloid fauna. — In: OLÓRIZ F. & F.J. RODRÍGUEZ-TOVAR (Eds), *Advancing Research in Living and Fossil Cephalopods*. Kluwer Academic / Plenum Publishers, New York: 365-379.
- HISTON K. (2002): A nautiloid assemblage from the Upper Silurian (Pídlí) of the Carnic Alps, Austria. — In: WYSE JACKSON P.N., PARKES M.A. & R. WOOD (Eds), *Studies in Palaeozoic Palaeontology and Biostratigraphy in Honour of Charles Hepworth Holland*. *Spec. Papers Palaeont.* **67**: 115-133.
- HISTON K. (2012a): Palaeoenvironmental and temporal significance of variably colored Paleozoic orthoconic nautiloid cephalopod accumulations. — In: FERRETTI A., HISTON K., McLAUGHLIN P. & C.E. BRETT (Eds), *Special Issue Time Specific Facies: the color and texture of biotic events*. *Palaeogeogr., Palaeoclimatol., Palaeoecol.* **367-368**: 193-208.
- HISTON K. (2012b): The Silurian nautiloid-bearing strata of the Cellon Section (Carnic Alps, Austria): colour variation related to events. — In: FERRETTI A., HISTON K., McLAUGHLIN P. & C.E. BRETT (Eds), *Special Issue Time Specific Facies: the color and texture of biotic events*. *Palaeogeogr., Palaeoclimatol., Palaeoecol.* **367-368**: 231-255.
- HISTON K., HUBMANN B. & F. MESSNER (2010): A preliminary study of the upper Silurian nautiloid cephalopods from the Eggenfeld section (Graz Paleozoic, Austria). — *Boll. Soc. Paleont. Italiana* **49**: 65-74.
- HOFSTETTER B., FUENZALIDA H. & G. CECIONI (1957): Chile. In: *Lexique Stratigraphique International*, Vol. 5. Centre de Recherche Sciences, Paris: 444 pp.
- HOLLAND C.H. (1998): The nautiloid cephalopod order Actinocerida in the British Silurian. — *Palaeontology* **41** (1): 183-192.
- HOLLAND C.H. (1999): The nautiloid cephalopod order Ascocerida in the British Silurian. — *Palaeontology* **42** (4): 683-689.
- HOLLAND C.H. (2000a): *Harrisoceras* and *Temperoceras* closely related cephalopod genera from the British Silurian. — *Boll. Soc. Paleont. Italiana* **39** (2): 117-122.
- HOLLAND C.H. (2000b): Silurian Cephalopods from the Pentland Hills. — *Scottish J. Geol.* **36**: 177-186.
- HOLLAND C.H. (2000c): Rare cephalopods from the Pridoli of England and Wales. — *Geol. J.* **35**: 25-31.
- HOLLAND C.H. (2002): Cephalopods from a borehole in the type Wenlock area. — *Proc. Geol. Ass.* **113**: 207-215.
- HOLLAND C.H. (2003): Some observations on bactritid cephalopods. — *Bull. Geosci.* **78**: 369-372.
- HOLLAND C.H. (2004): The curious case of '*Orthoceras primaevum*' from the Silurian of Wales. — *Proc. Geol. Ass.* **115**: 235-238.
- HOLLAND C.H. (2007): Cyrtoconic nautiloid cephalopods from the British Silurian. — *Proc. Geol. Ass.* **118**: 365-373.
- HOLLAND C.H. (2010): Coiled nautiloid cephalopods from the British Silurian. — *Proc. Geol. Ass.* **121**: 13-23.
- HOLLAND C.H. & S. STRIDSBERG (2004): Specific representation of the Silurian cephalopod genus *Phragmoceras* in Gotland and Britain. — *GFF* **126**: 301-310.
- HOLM G. (1891): Om mynningen hos *Lituities* BREYN. — *Geol. Fören. Stockholm Förhandl.* **13**: 736-774.
- HOUSE M.R. (1978): Devonian ammonoids from the Appalachians and their bearing on international zonation and correlation. — *Spec. Papers Palaeont.* **21**: 1-70.
- HOWARTH M.K. (1965): Cretaceous ammonites and nautiloids from Angola. — *Bull. British Mus. (Nat. Hist.), Geol.* **10**: 335-412.
- HYATT A. (1883-1884): Genera of fossil cephalopods. — *Proc. Boston Soc. Nat. Hist.* **22**: 253-338.
- HYATT A. (1891): Carboniferous cephalopods. — *Rep. Geol. Survey Texas, Austin* **2**: 329-356.
- HYATT A. (1893): Carboniferous cephalopods, paper 2. — *Rep. Geol. Survey Texas, Austin* **4**: 377-474.
- HYATT A. (1894): Phylogeny of an acquired characteristic. — *Proc. American Philos. Soc.* **22**: 349-647.
- JAANUSSON V. (1963): Lower and Middle Viruan (Middle Ordovician) of the Siljan District. — *Bull. Geol. Inst. Univ. Uppsala* **42**: 1-40.
- JAANUSSON V. & H. MUTVEI (1953): Stratigraphie und Lithologie der unterordovizischen Platyrus-Stufe in Siljan-Gebiet, Dalarna. — *Bull. Geol. Inst. Univ. Uppsala* **35**: 7-34.
- JAEGER H. (1975): Die Graptolithenführung im Silur/Devon des Cellon-Profiles (Karnische Alpen). — *Carinthia II* **165**: 111-126.
- JOHNSON M.E. (2010): Tracking Silurian eustasy: Alignment of empirical evidence or pursuit of deductive reasoning? — *Palaeogeogr., Palaeoclimatol., Palaeoecol.* **296**: 276-284.
- KING A.H. (1990): Lower and Middle Ordovician Cephalopoda of Baltoscandia. Unpubl. Ph.D. Thesis, University of Wales, Swansea.
- KING A.H. (1999): A review of Volkhovian and Kundan (Arenig-Llanvirn) nautiloids from Sweden. — In: OLÓRIZ F. & F.J. RODRÍGUEZ-TOVAR (Eds), *Advancing Research on Living and Fossil Cephalopods*. Kluwer Academic / Plenum Publishers. New York: 137-159.
- KING A.H. (2011): Fossil nautiloids from the Upper Lias (Toarcian) 'Junction Bed' of the Ilminster area, Somerset. — *Proc. Somerset Archaeol. Nat. Hist. Soc.* **154**: 249-258.
- KLEFFNER M.A. (1989): A conodont-based Silurian chronostratigraphy. — *Geol. Soc. America Bull.* **101**: 904-912.
- KLEFFNER M.A. (1995): A conodont- and graptolite-based Silurian chronostratigraphy. — In: MANN K.O. & H.R. LANE (Eds): *Graphic Correlation. Soc. Sed. Geol. (Spec. Publ.)* **53**: 159-176.
- KLUG C. (2002): Quantitative stratigraphy and taxonomy of late Emsian and Eifelian ammonoids of the eastern Anti-Atlas (Morocco). — *Cour. Forschungsinst. Senckenberg* **238**: 1-109.
- KLUG C., KRÖGER B., KIESSLING W., MULLINS G.L., SERVAIS T., FRYDA J., KORN D. & S. TURNER (2010): The Devonian nekton revolution. — *Lethaia* **43**: 465-477.

- KOLEBABA I. (1975): *Caliceras* n. gen. and ontogeny of *C. capillosum* (BARRANDE) (Nautiloidea, Michelinoceratida). — *Casopis Mineral. Geol.* **20**: 377-391.
- KOLEBABA I. (1977): New information on longitudinally sculptured orthoceroids. — *Casopis Mineral. Geol.* **22**: 125-138.
- KOLEBABA I. (1999): Sipho-cameral structures in some Silurian cephalopods from the Barrandian area (Bohemia). — *Acta Mus. Nat. Pragae, Ser. B: Hist. Nat.* **55** (1-2): 1-15.
- KOLEBABA I. (2002): A contribution to the theory of the cameral mantle in some Silurian Nautiloidea; Mollusca, Cephalopoda. — *Bull. Czech Geol. Survey* **77** (3): 183-186.
- KORN D. (1996): Revision of the Rhenish Late Visean goniatite stratigraphy. — *Ann. Soc. géol. Belgique* **117**: 129-136.
- KŘÍŽ J. (1998): Recurrent Silurian-lowest Devonian cephalopod limestones of Gondwanan Europe and Perunica. — In: LANDING E. & M.E. JOHNSON (Eds), *Silurian Cycles Linkages of Dynamic Stratigraphy with Atmospheric, Oceanic and Tectonic Changes*. James HALL Centennial Volume. New York State Mus. Bull. **491**: 183-198.
- KŘÍŽ J., DEGARDIN J.M., FERRETTI A., HANSCH W., GUTIÉRREZ-MARCO J.C., PARIS F., PICARRA D'ALMEIDA J.M., ROBARDET M., SCHÖNLAUB H.P. & E. SERPAGLI (2003): Silurian stratigraphy and paleogeography of Gondwanan and Perunican Europe. — In: LANDING E. & M.E. JOHNSON (Eds), *Silurian Lands and Seas: Paleogeography Outside of Laurentia*. New York State Mus. Bull. **493**: 105-178.
- KRÖGER B. (2006): Early growth-stages and classification of orthocerid cephalopods of the Darriwilian (Middle Ordovician) of Baltoscandia. — *Lethaia* **39**: 129-139.
- KRÖGER B. (2008a): Nautiloids before and during the origin of ammonoids in a Siluro-Devonian section in the Tafilalet, Anti-Atlas, Morocco. — *Spec. Papers Palaeont.* **79**: 1-110.
- KRÖGER B. (2008b): A new genus of middle Tremadocian orthoceratoids and the Early Ordovician origin of orthoceratoid cephalopods. — *Acta Palaeont. Polonica* **53**: 745-749.
- KRÖGER B. (2013): The cephalopods of the Boda Limestone, Late Ordovician, of Dalarna, Sweden. — *European J. Taxon.* **41**: 1-110. <http://dx.doi.org/10.5852/ejt.2013.41>
- KRÖGER B. & D.H. EVANS (2011): Review and palaeoecological analysis of the late Tremadocian-early Floian (Early Ordovician) cephalopod fauna of the Montagne Noire, France. — *Fossil Rec.* **14**: 5-34. doi:10.1002/mmng.201000013
- KRÖGER B. & E. LANDING (2008): Onset of the Ordovician cephalopod radiation – evidence from the Rochdale Formation (middle Early Ordovician, Stairsian) in eastern New York. — *Geol. Mag.* **145**: 1-31. doi:10.1017/S0016756808004585
- KRÖGER B. & B. LEFEBVRE (2012): Palaeogeography and palaeoecology of early Floian (Early Ordovician) cephalopods from the Upper Fezouata Formation, Anti-Atlas, Morocco. — *Fossil Rec.* **15**: 61-75. doi:10.1002/mmng.201200004
- KRÖGER B., SERVAIS T. & Y. ZHANG (2009): The origin and initial rise of pelagic cephalopods in the Ordovician. — *PLoS ONE* **4** (9): e7262. doi:10.1371/journal.pone.0007262
- KUMMEL B. (1953): American Triassic coiled nautiloids. — *U.S. Geol. Survey Prof. Paper* **250**: 1-104.
- KUMMEL B. (1956): Post-Triassic nautiloid genera. — *Bull. Harvard Mus. Comp. Zool.* **114** (7): 319-494
- KUMMEL B. (1964): Nautilida. — In: MOORE R.C. (Ed.), *Treatise on Invertebrate Paleontology, Part K. Mollusca 3*. Geological Society of America and University of Kansas Press, New York and Lawrence: K383-456.
- LAI C.-G. (1982): Ordovician cephalopods from Xainza, Xizang (Tibet). — *Acta Palaeont. Sinica* **21** (5): 553-559. (In Chinese with English summary and description of new species).
- LAI C.-G. & M.-Q. WANG (1981): Nautiloidea. — In: *Palaeontological Atlas of North West China: Xinjiang Uygur Autonomous Region, Vol. 1. Late Proterozoic – Early Palaeozoic*. Geological Publishing House, Beijing: 115-134. [In Chinese]
- MANDA Š. (1996): *Cyrtograptus lundgreni* Biozone in the southwestern part of the Svatý Jan Volcanic Centre (Wenlock, Prague Basin, Bohemia). — *Vstník eského geol. ústavu* **71** (4): 369-374.
- MANDA Š. (2008): Palaeoecology and palaeogeographic relations of the Silurian phragmoceratids (Nautiloidea, Cephalopoda) of the Prague Basin (Bohemia). — *Bull. Geosci.* **83** (1): 39-62.
- MANDA Š. & J. FRYDA (2010): Silurian-Devonian boundary events and their influence on cephalopod egg size during mass extinctions. — *Bull. Geosci.* **85**: 513-540.
- MANDA Š. & J. KŘÍŽ (2006): Environmental and biotic changes in subtropical isolated carbonate platforms during the Late Silurian Kozłowski Event, Prague Basin. — *GFF* **128**: 161-168.
- MANDA Š. & J. KŘÍŽ (2007): New cephalopod limestone horizon in the Ludlow (Gorstian, lower *L. scanicus* Biozone) of the Prague Basin (Bohemia, Perunica). — *Boll. Soc. Paleont. Italiana* **46** (1): 33-45.
- MANDA Š. & V. TUREK (2009): Revision of the Pragian Rutoceratoidea Hyatt, 1884 (Nautiloidea, Oncocerida) from the Prague Basin. — *Bull. Geosci.* **84**: 127-148.
- MAPES R.H., NIKO S., FRYDA J. & A. NÜTZEL (2007): A newly hatched coiled nautiloid from the Permian of Italy. — *J. Paleont.* **81**: 1118-1121.
- MILLER A.K. & W.M. FURNISH (1938): Lower Mississippian nautiloid cephalopods of Missouri. — In: *Stratigraphy and Paleontology of the Lower Mississippian of Missouri. Pt. 2. Missouri Univ. Studies* **13** (4): 148-178.
- MILLER A.K. & H.F. GARNER (1953): Lower Mississippian cephalopods of Michigan. Pt. 2. Coiled Nautiloids. — *Contr. Mus. Paleont., Univ. Michigan* **11**: 111-151.
- MILLER A.K. & J.B. OWEN (1934): Cherokee nautiloids of the northern mid-continent region. — *Stud. Nat. Hist. Iowa Univ.* **16**: 187-272.
- MILLER A.K. & A.G. UNKLESBAY (1942): Permian nautiloids from western United States. — *J. Paleont.* **16**: 719-738.
- MILLER A.K. & W.L. YOUNGQUIST (1949): American Permian nautiloids. — *Geol. Soc. America Mem.* **41**: 1-218.
- MILLER A.K., DUNBAR C.O. & G.E. CONDRA (1933): The nautiloid cephalopods of the Pennsylvanian system in the Mid-continent region. — *Bull. Geol. Survey Nebraska, Ser. 2* **9**: 1-240.
- MILLER A.K., LANE J.H. & A.G. UNKLESBAY (1947): A nautiloid cephalopod fauna from the Pennsylvanian Winterset Limestone of Jackson County, Missouri. — *Contr. Paleont. Kansas, Lawrence, Mollusca* **2**: 1-11.
- NIELSEN S.N. & C. SALAZAR (2011): *Eutrephoceras subplicatum* (STEINMANN, 1895) is a junior synonym of *Eutrephoceras dorbignyanum* (FORBES in DARWIN, 1846) (Cephalopoda, Nautiloidea) from the Maastrichtian Quiriquina Formation of Chile. — *Cretaceous Res.* **32**: 833-840.
- NIKO S. (1990): Early Carboniferous Visean cephalopods from the Hikoroichi Formation, southern Kitakami Mountains. — *Trans. Proc. Palaeont. Soc. Japan* **159**: 554-561.

- NIKO S., NISHIDA T. & Y. KYUMA (1987): Middle Carboniferous Orthocerataceae and Pseudorthocerataceae from the Akiyoshi limestone. — Trans. Proc. Palaeont. Soc. Japan **148**: 335-345.
- NOETLING F. (1884): Beiträge zur Kenntniss der Cephalopoden aus Silurgeschieben der Provinz Ost-Preussen. — Jahrb. Königl. Preussischen Geol. Landesanst. Bergakad. Berlin **1883**: 101-135.
- OPPEL A. (1865): Geognostische Studien in dem Ardèche Dept. — Paläont. Mitt. Mus. Königl. Bayer. Staates **5**: 305-322.
- PAGE K.N. (2009): High resolution ammonite stratigraphy of the Charmouth Mudstone Formation (Lower Jurassic: Sinemurian-Lower Pliensbachian) in south-west England, UK. — Vol. Jurassica **7**: 19-29.
- PHILLIPS D. (1982): Catalogue of the Type and Figured Specimens of Fossil Cephalopoda (excluding Mesozoic Ammonoidea) in the British Museum (Natural History). — British Museum (Natural History), London: 94 pp.
- PHILLIPS D. (1985): The nautiloid *Brachycycloceras* in the Upper Carboniferous of Britain. — Palaeontology **28**: 235-242.
- PIA J. (1914): Untersuchungen über die liassischen Nautiloidea. — Beitr. Paläont. Geol. Österr.-Ungarns u. d. Orients **27**: 19-86.
- POSENATO R. & H. PREINOTH (2004): Orizzonti a nautiloidi e brachiopodi della Formazione a Bellerophon (Permiano Superiore) in Val Gardena (Dolomiti). — Geo.Alp **1**: 71-85.
- POSENATO R. (2010): Marine biotic events in the Lopingian succession and latest Permian extinction in the Southern Alps (Italy). — Geol. J. **45**: 195-215.
- PRIEWALDER H. (1997): The distribution of the chitinozoans in the Cellon section (Hirnantian-lower Lochkovian) – a preliminary report. — In: SCHÖNLAUB H.P. (Ed.), IGCP – 421 Inaugural Meeting Vienna, Guidebook. Ber. Geol. Bundesanst. **40**: 74-85.
- QI D.-L. (1980): Ordovician cephalopods from Wuwei of Anhui and their stratigraphical significance. — Acta Palaeont. Sinica **19**: 245-262. [In Chinese with English abstract p. 273].
- REMELE A. (1880): Ueber einige neue oder seltene Versteinerungen aus silurischen Diluvialgeschieben der Gegend von Eberswalde. — Festschrift für die 50jährige Jubelfeier der Forstakademie Eberswalde, Eberswalde: 179-252.
- REMELE A. (1881): Ueber einige gekrümmte untersilurische Cephalopoden. — Zeitschr. Deutschen Geol. Gesell. **34**: 1-68.
- REYMENT R.A. (1958): Some factors in the distribution of fossil cephalopods. — Stockholm Contrib. Geol. **1**: 97-184.
- RISTEDT H. (1968): Zur Revision der Orthoceratidae. — Akad. Wiss. u. Lit., Abhandl. math.-naturwiss. Kl. **4**: 213-297.
- RISTEDT H. (1969): Orthoceren als Leitfossilien des Silurs. — Carinthia II **27**: 25-28.
- RISTEDT H. (1981): Bactriten aus dem Obersilur Böhmens. — Mitt. Geol.-Paläontol. Inst. Univ. Hamburg **51**: 23-26.
- RULLEAU L. (2008): Les Nautilus de Lias et du Dogger de la région Lyonnaise: 151 pp.
- SEGERSTROM K. (1960): Cuadrángulo Quebrada Paipote, Provincia de Atacama. — Instituto de Investigaciones Geológicas, Carta Geologica de Chile **2**: 1-35.
- SERPAGLI E. & M. GNOLI (1977): Upper Silurian Cephalopods from Southwestern Sardinia. — Boll. Soc. Paleont. Italiana **16** (2): 153-196.
- SERVENTI P. & R. FEIST (2009): Silurian nautiloid cephalopods from the Cabrières area (Montagne Noire, France): a preliminary report. — In: CORRIGA M.G. & S. PIRAS (Eds), Time and Life in the Silurian: A Multidisciplinary Approach. Abstracts. Rendiconti Soc. Paleont. Italiana **3** (3): 341.
- SERVENTI P. & M. GNOLI (2000): Nouvi ritrovamenti di Cefalopodi nautiloidi nelle Alpi Carniche. — Giorn. Geol., ser. 3 **62**: 9-14.
- SERVENTI P., CORRADINI C., SIMONETTO L. & M. PONDRELLI (2006): Silurian nautiloid cephalopods of the Museo Friulano di Storia Naturale: Family Orthoceratidae. — Gortania **28**: 29-57.
- SERVENTI P., GNOLI M., & L. SIMONETTO (2010): Actinoceratid Cephalopods from the Silurian of the Carnic Alps (Italian side). — Boll. Soc. Paleont. Italiana **49** (1): 75-81.
- SHIMANSKY V.N. (1948): Some new orthoceracones from Artinskian deposits of the Southern Urals. — Doklady Akad. Nauk SSSR, Leningrad **60**: 119-121. [In Russian]
- SHIMANSKY V.N. (1957): Carboniferous Oncoceratida. — Doklady Akad. Nauk SSSR, Leningrad **112**: 530-532. [In Russian]
- SHIMANSKY V.N. (1967): Carboniferous Nautiloidea. — Akad. Nauk. SSSR, Paleont. Inst. Tr. **115**: 1-258. [In Russian]
- SHIMANSKY V.N. (1968): Carboniferous Orthoceratida, Oncoceratida, Actinoceratida, Bactritida. — Akad. Nauk. SSSR, Paleont. Inst. Tr. **117**: 1-148. [In Russian]
- SMITH W. (1816): Strata Identified by Organised Fossils, Containing Prints on Coloured Paper of the Most Characteristic Specimens in Each Stratum. — W. Arding, London.
- SOBOLEV E.B. (1994): Stratigraphic range of Triassic boreal Nautiloidea. — In: GUÉX J. & A. BAUD (Eds), Recent Developments on Triassic Stratigraphy. Université de Lausanne, Institut de Géologie et Paléontologie, Lausanne: 127-138.
- SOWERBY J. (1815-18): The Mineral Conchology of Great Britain. Volume II. — Longman & Co. and Sherwood & Co., London: 239 pp.
- SPATH L.F. (1927): Revision of the Jurassic cephalopod fauna of the Kachh (Cutch). — Mem. Geol. Survey India (Palaeont. Indica), New Ser. **9** (2): 1-84.
- SPATH L.F. (1953): The Upper Cretaceous cephalopod fauna of Graham Land. — Falkland Islands Depend. Survey Sci. Rep. **3**: 2-58.
- STEINMANN G. (1895): Die Cephalopoden der Quiriquina-Schichten. — N. Jahrb. Mineral. Geol. Paläont. **10**: 64-68.
- STINNESBECK W. (1986): Zu den faunistischen und palökologischen Verhältnissen in der Quiriquina Formation (Maastichtium) Zentral-Chiles. — Palaeontographica A **194**: 99-237.
- STRIDSBERG S. (1985): Silurian Oncocerid cephalopods from Gotland. — Fossils and Strata **18**: 1-65.
- STRIDSBERG S. & V. TUREK (1997): A revision of the Silurian nautiloid genus *Ophioceras* BARRANDE. — GFF **19**: 21-36.
- STURGEON M.T. (1946): Allegheny fossil invertebrates from Eastern Ohio Nautiloidea. — J. Paleont. **20**: 8-37.
- STURGEON M.T. & A.K. MILLER (1948): Some additional cephalopods from the Pennsylvanian of Ohio. — J. Paleont. **22**: 75-80.
- STURGEON M.T., WINDLE D.L., MAPES R.H. & R.D. HOARE (1997): Nautiloid and bactritoid cephalopods. — Ohio Div. Geol. Survey Bull. **71**: 1-191.

- SWEET W.C. (1958): The Middle Ordovician of the Oslo Region, Norway. 10. Nautiloid cephalopods. — Norsk Geol. Tidsskr. **38**: 1-176.
- TEICHERT C. & B.F. GLENISTER (1954): Fossil nautiloid faunas from Australia. — J. Paleont. **26**: 730-752.
- TEICHERT C. & B. KUMMEL (1973): Nautiloids from the Julfa Beds, Upper Permian, northwest Iran. — Bull. Mus. Comp. Zool. **144**: 409-434.
- TEICHERT C., KUMMEL B., SWEET W.C., STENZEL H.B., FURNISH W.M., GLENISTER B.F., ERBEN H.K., MOORE R.C. & D.E. NODINE ZELLER (1964): Treatise on Invertebrate Paleontology. Part K. Mollusca 3. Cephalopoda – General Features. Endoceratoidea – Actinoceratoidea – Nautiloidea – Bactritoidea. — University of Kansas Press and Geological Society of America, Lawrence, Kansas: 519 pp.
- TERQUEM O. (1855): Paléontologie de l'étage inférieur de la Formation Liassique de la Province de Luxembourg (Grand-Duché) et de Hettange, Département de la Moselle. — Mém. Soc. Géol. France **5**: 1-125.
- TINTANT H. (1984): Contribution a la connaissance des nautilices jurassique 1. – Le sous-genre *Cenoceras* HYATT dans le Lias du Sud-Est de la France. — Geol. France **1-2**: 29-66.
- TINTANT H. (1987): Les nautilices du Jurassique d'Arabie Saoudite. — Geobios, Mém. Spec., **9**: 67-159.
- TUREK V. (1975): Genus *Kosovoceras* gen. n. in the Silurian of Central Bohemia (Nautiloidea). — Sbor. Geol. Ved., Paleont. **17**: 7-44.
- TUREK V. (2008): *Boionautilus* gen. nov. from the Silurian of Europe and North Africa (Nautiloidea, Tarphycerida). — Bull. Geosci. **83** (2): 141-152.
- TURNER J.S. (1937): The faunal succession in the Carboniferous Limestone near Cork. — Proc. R. Irish Acad. **43B**: 193-209.
- TURNER J.S. (1939): The upper Devonian and lower Carboniferous of the Cork district. — In: The Geology of Southeast Ireland, Together with Parts of Limerick, Clare and Galway. Proc. Geol. Ass. **3**: 287-351.
- TURNER J.S. (1950): The Carboniferous limestone in Co. Dublin south of the River Liffey. — Sci. Proc. R. Dublin Soc. **25**: 169-192.
- TURNER J.S. (1951): On the Carboniferous nautiloids: *Orthocera gigantea* J. SOWERBY and allied forms. — Trans. R. Soc. Edinburgh **62** (1): 169-190.
- TURNER J.S. (1953): The nautiloids named in MARTIN'S 'Petreficata Derbiensia, 1809'. — Ann. Mag. Nat. Hist., ser. 12 **6**: 689-692.
- TURNER J.S. (1954): On the Carboniferous nautiloids: Some Middle Visian species from the Isle of Man. — Liverpool and Manchester Geol. J. **1**: 298-325.
- TURNER J.S. (1965): On the Carboniferous nautiloids: *Nautilus quadratus* FLEMING and certain other coiled nautiloids. — Proc. Leeds Phil. Lit. Soc., sci. sect. **9**: 223-256.
- TURNER J.S. (1966): On the Carboniferous nautiloids: The Phacoceratidae. — Proc. Leeds Phil. Lit. Soc., sci. sect. **10**: 1-15.
- ULRICH E.O., FOERSTE A.F., MILLER A.K. & A.G. UNKLESBAY (1944): Ozarkian and Canadian cephalopods. Part III: Longicones and summary. — Geol. Soc. America Spec. Papers **58**: 1-226.
- WALLISER O.H. (1964): Conodonten des Silurs. — Abh. Hessischen L.-Amtes Bodenforsch. **41**: 1-106.
- WENZEL B.C. (1997): Isotopenstratigraphische Untersuchungen an silurischen Abfolgen und deren paläoozeanische Interpretation. — Erlanger Geol. Abh. **129**: 1-117.
- WILSON A.E. (1961): Cephalopoda of the Ottawa Formation of the Ottawa St. Lawrence Lowland. — Geol. Survey Canada, Bull. **67**: 1-106.
- YU C.C. (1930): The Ordovician cephalopods of central China. — Palaeont. Sinica, B **1**: 1-71.
- YUN C-S. (1999): Ordovician cephalopods from the Maggor Formation of Korea. — Paleont. Res. **3**: 202-221.
- YUN C-S. (2002): Faunal and biogeographical characteristics of the Ordovician cephalopods from Korea. — Abh. Geol. Bundesanst. **57**: 555-569.
- ZHURAVLEVA F.A. (1974): Devonian nautiloids. Orders Oncocerida, Tarphycerida and Nautilida. — Trudy Paleont. Inst. **142**: 1-160.

David H. EVANS
Natural England
Unex House
Bourges Boulevard
Peterborough, PE1 1UA, UK
E-Mail: david.evans@naturalengland.org.uk

Andy H. KING
Geckoella Ltd
Suite 323
7 Bridge Street
Taunton, Somerset, TA1 1TD, UK
E-Mail: asia.nautilus@gmail.com

Kathleen HISTON
Via Mazzini 4
21039 Valganna
Varese, Italy
E-Mail: hiscat@interfree.it

Marcela CICHOWOLSKI
Área de Paleontología
Departamento de Ciencias Geológicas
Universidad de Buenos Aires
Ciudad Universitaria, Pab. II
Intendente Güiraldes 2160 C1428EGA
Buenos Aires, Argentina
E-Mail: mcicho@gl.fcen.uba.ar